

## Stratégie en 5 points

### Pour conserver des liens avec les proches de personnes hospitalisées aux soins intensifs en raison de la COVID-19

Tiré de : Azoulay, E., Kentish-Barnes, N. (2020). A 5-point strategy for improved connection with relatives of critically ill patients with COVID-19. *The Lancet*. [https://doi.org/10.1016/S2213-2600\(20\)30223-X](https://doi.org/10.1016/S2213-2600(20)30223-X)

Traduction libre par Catherine Wilhelmy, Réseau Planetree francophone.

- 1 **STIMULER LA VISITE DES PROCHES** en les restreignant aux plus intimes, asymptomatiques, et en mesure d'appliquer de façon optimale les mesures de protection.
- 2 **STANDARDISER L'INFORMATION ÉCRITE DESTINÉE AUX PROCHES.** À la suite d'une discussion entre les proches et les cliniciens, un écrit est envoyé aux proches et reprend les principaux éléments de la discussion qui s'est déroulée entre eux.
- 3 **PLANIFIER UN APPEL DE ROUTINE AVEC LES PROCHES,** afin de maintenir une continuité dans la communication. Les appels pourraient être effectués par des étudiants en médecine, des médecins (autres que ceux de l'équipe des soins intensifs), d'autres professionnels du réseau de la santé ou des bénévoles [formés]. En plus de ces appels, les proches sont invités à contacter le médecin ou l'infirmière une fois par jour afin d'obtenir les informations de base. De l'information plus détaillée est fournie à l'occasion d'une rencontre en personne lorsque les proches visitent le patient. Pour les patients conscients, les téléphones intelligents & tablettes électroniques peuvent contribuer à des visites virtuelles.
- 4 **DEMEURER EN CONTACT AVEC LE PATIENT.** Encourager les proches à utiliser les moyens qui répondent aux besoins du patient d'être en contact avec les siens (conférences par internet avec la famille, journal de famille, dessins, messages textes, groupes sur des réseaux sociaux).
- 5 **CHANGER D'APPROCHE POUR LES PERSONNES EN FIN DE VIE** afin d'éviter de priver des proches de leurs derniers moments ensemble. Par exemple, si nécessaire, une rencontre de famille virtuelle devrait être organisée afin de répondre au besoin de la famille de se préparer au deuil. Un proche devrait pouvoir rester dans la chambre autant que possible. Toutefois, la sécurité des proches constitue une priorité, et l'équipe de soin doit pouvoir fournir à la personne qui visite, la formation et l'équipement nécessaire à sa présence.

## A 5-point strategy for improved connection with relatives of critically ill patients with COVID-19

The coronavirus disease 2019 (COVID-19) pandemic has spread throughout the world, leading hospitals to expand their critical-care capacity.<sup>1</sup> Logistics in times of surging demand are challenging. Health-care providers are overwhelmed by the relentless workload and tend to focus on the patients, as they have little time for family-centred care.

In many places, a nearly complete lockdown combined with stringent social distancing measures have been put in place in an attempt to curb the spread of the COVID-19 pandemic. Hospital visits are prohibited to ensure that relatives do not contaminate other family members, patients, or health-care professionals. However, the burden on intensive care unit (ICU) relatives of patients with COVID-19 is particularly heavy.<sup>2</sup> The lockdown imposed by many governments can result in confusion, stress, frustration, anger, communication gaps, and post-traumatic stress-related symptoms.<sup>3</sup> People are more exposed than usual to the media, whose messages often cause additional distress. Moreover, studies have shown that family members of survivors of acute respiratory distress syndrome have higher rates of post-ICU burden than the patients themselves.<sup>4,5</sup> Masks and other protective equipment hamper the ability of health-care professionals to express empathy and provide fine-tuned communication. As a result, patients could be in a suboptimal position to make decisions, and families might feel frustrated. Finally,

post-traumatic stress disorder is significantly more common in bereaved relatives when the patients were seen with shortness of breath before death.<sup>6</sup>

We suggest a 5-point strategy to foster a positive connection between patients, relatives, and health-care professionals during the COVID-19 pandemic (panel). Studies are needed to identify the specific communication needs of relatives of critically ill patients with COVID-19. Moreover, whether these family members suffer a greater post-ICU burden than relatives of other ICU patients with acute respiratory failure (for instance, due to community-acquired pneumonia or influenza) deserves further exploration. This 5-S strategy to maintain the connection with families and patients with COVID-19 aims primarily to improve communication with highly vulnerable relatives. Post-traumatic stress disorder has been shown to be associated with suboptimal communication, particularly in relatives of patients who died in the ICU.<sup>8</sup> In this setting, the quality of interactions between family and health-care professionals is a major determinant of complicated grief.<sup>6</sup>

We declare no competing interests.

\**Elie Azoulay, Nancy Kentish-Barnes*  
 elie.azoulay@aphp.fr

Médecine Intensive et Réanimation, Hôpital Saint-Louis AP-HP, Paris University, Paris 75010, France

- 1 Phua J, Weng L, Ling L, et al. Intensive care management of coronavirus disease 2019 (COVID-19): challenges and recommendations. *Lancet Respir Med* 2020; published online April 6. [https://doi.org/10.1016/S2213-2600\(20\)30161-2](https://doi.org/10.1016/S2213-2600(20)30161-2).
- 2 Holmes EA, O'Connor RC, Perry VH, et al. Multidisciplinary research priorities for the COVID-19 pandemic: a call for action for mental health science. *Lancet Psychiatry* 2020; published online April 15. [https://doi.org/10.1016/S2215-0366\(20\)30168-1](https://doi.org/10.1016/S2215-0366(20)30168-1).
- 3 Brooks SK, Webster RK, Smith LE, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet* 2020; **395**: 912–20.

- 4 Dinglas VD, Chessare CM, Davis WE, et al. Perspectives of survivors, families and researchers on key outcomes for research in acute respiratory failure. *Thorax* 2018; **73**: 7–12.
- 5 Sanfilippo F, Ippolito M, Santonocito C, et al. Long-term functional and psychological recovery in a population of acute respiratory distress syndrome patients treated with VV-ECMO and in their caregivers. *Minerva Anestesiol* 2019; **85**: 971–80.
- 6 Kentish-Barnes N, Chaize M, Seegers V, et al. Complicated grief after death of a relative in the intensive care unit. *Eur Respir J* 2015; **45**: 1341–52.
- 7 Lautrette A, Darmon M, Megarbane B, et al. A communication strategy and brochure for relatives of patients dying in the ICU. *N Engl J Med* 2007; **356**: 469–78.
- 8 Azoulay E, Pochard F, Kentish-Barnes N, et al. Risk of post-traumatic stress symptoms in family members of intensive care unit patients. *Am J Respir Crit Care Med* 2005; **171**: 987–94.



*Lancet Respir Med* 2020

Published Online

May 4, 2020

[https://doi.org/10.1016/S2213-2600\(20\)30223-X](https://doi.org/10.1016/S2213-2600(20)30223-X)

### Panel: A 5-point (5S) strategy to maintain the connection with relatives of critically ill patients with COVID-19

- 1 Stimulate family visits by restricting them to the closest relative who is asymptomatic and able to apply optimal personal protective measures.
- 2 Standardised written information for the relatives. This letter is sent to the relatives but also discussed during the first remote call between the health-care professionals and the family.
- 3 Schedule routine telephone calls with the relatives to maintain continuity of communication. The calls could be made by a medical student, a non-ICU physician, another health-care professional, or a volunteer. In addition to these calls, family members are invited to call the physician and bedside nurse once a day to receive basic information. More detailed information is to be provided during a face-to-face meeting when the relative visits the patient. For conscious patients, smartphones and digital tablets can allow video calls or virtual ICU visits.
- 4 Stay in touch by encouraging the family to find ways to improve the link with the relative (web-based remote family conferences, diaries, drawings, text messages, and media groups).
- 5 Switch to a different approach in end-of-life situations to avoid depriving family members of the opportunity to say goodbye to a relative. For example, family conferences should be organised, remotely if needed, to meet family needs and prepare for the bereavement.<sup>7</sup> The relative should be allowed to stay in the room as much as possible. However, safety of the relative is a primary concern, and the health-care professionals must provide the relative with training in the use of personal protective equipment.